

Housekeeping Rules

- Place questions in the “Q&A” chat feature – questions will be answered at the end of the presentation.
- Place technical issues in the chat box.

Zoom Webinar
Enter Full Screen

Erik Casas

Chat Box

Q&A
Chat
Feature

Join Audio Chat Raise Hand Q&A Leave

Strategies for integrating suicide risk assessment and management into evidence-based treatments for PTSD

September 8, 2022

Presented By:

Brooke Fina, LCSW, BCD

Associate Professor-Research

Department of Psychiatry and Behavioral Sciences, School of Medicine

University of Texas Health Science Center at San Antonio

5


PTSD & Suicidal Thoughts and Behaviors in Veterans

- *Traum a exposure* increases risk for suicide (e.g., childhood abuse, military sexual traum a) (Afifi et al., 2008; Ystgaard et al., 2004).
- In a national sam ple of veterans, PTSD increased the odds of SI (OR=9.7) and suicide attempts (OR=11.8) (Wisco et al., 2014).
- Iraq and Afghanistan Veterans with PTSD and 2 or more com orbid mental disorders are 5.7x m ore likely to have suicidal thoughts com pared to veterans with PTSD (Jakupcaket et al., 2009).

QUESTIONS?


Thank you!

fina@uthscsa.edu




Q&A

- Please type your questions in the Q&A feature at the bottom of the screen.



Considerations for Suicide Risk Management in PTSD Tx: Treatment Phase



PTSD	Suicide Risk
PTSD Checklist for DSM-5 (PCL-5)	Depressive Symptom Inventory Suicidality Subscale (DSI-SS)
Patient Health Questionnaire-9 (PHQ-9)	Suicide Cognitions Scale-Short Form (SCS-S)
	Pre-Tx: Develop initial CRP as a part of treatment planning